A B

Figure 2. A, Preoperative view of a 52-year-old woman.B, Postoperative view 1 week after undergoing WAL.

A B

Figure 3. A, Preoperative view of a 60-year-old-man.B, Postoperative view 2 days after undergoing WAL.

4. Considerably less intraoperative swelling allows the surgeon to realize the target result with greater precision.
5. Fine shaping of small deposits with precision is also possible. This enables surgeons to accommodate the increasing patient demands for contouring.

It is notable that patient safety has increased considerably, even in extensive procedures. The limitations and drawbacks of other in“ltration methods are reduced or eliminated. This new method of simultaneous ”uid in and out allows the surgeon and patient new consistent levels of comfort and security.■

A B

Figure 4. A, Preoperative view of a 51-year-old woman.B, Postoperative view 1 week after undergoing WAL.

A B

Figure 5. A, Preoperative view of a 51-year-old woman.B, Postoperative view 1 day after undergoing WAL.

Dr. Man received no financial support from and has no commercial interest in the manufacture of the products discussed in this article.

Dr. Meyer is an unpaid consultant to human med.

References

1. Klein J.A. The tumescent technique for liposuction surgery. *J Am Acad Cosmetic Surg* 1987;4:263-267.
2. Klein JA. The tumescent technique: anesthesia and modi"ed liposuction technique. *Dermatol Clin* 1990;8:424-437.
3. Zocchi ML. Ultrasonic-assisted lipoplasty. *Aesthetic Plast Reconstr Surg* 1996;23:575-598.
4. Gasparotti M. Super"cial liposuction: a new application for the technique for aged and "accid Skin. *Aesthetic Plast Surg* 1992;16:141-153.
5. Neira R, Ortiz-Neira C. Low level laser-assisted liposculpture: clinical report of 700 cases. *Aesthetic Surg J* 2002;22:451-455.

6. Tucker GT. Local anaesthetic drugs „ mode of action and pharmacokinetics. *Anaesthesia* 1990;38:983-1010.

7. McCaughey W. Adverse effects of local anesthetics. *Drug Safety* 1992;49:126-132.

8. Lindenblatt N, Belusa L, Teifenbach B, Schareck W, Olbrisch RR. Prilocaine plasma levels and methemoglobinemia in patients undergoing tumescent liposuction involving less than 2000 ml. *Aesthetic Plast Surg* 2004;28:435-440.

9. Housman T, Lawrence N, Mellen BG, Beorge MN, Filippo JS, Cerveny KA, et al. The safety of liposuction: results of a national survey. *Dermatologic Surg* 2002;28:971-978.

10. Weinberg GL, Laurito CE, Geldner P, Pygon BH, Burton BK. Malignant ventricular dysrhythmias in a patient with isovaleric academia receiving general and local anesthesia for suction lipectomy. *J Clin Anesth* 1997;9:668-670.

11. Howland MA. Pharmacokinetics and toxicokinetics. In: Goldfrank LR, editor. Goldfranks•s toxicologic emergencies. 6th ed. Stanford, CT: Appleton & Lange; 1998. p. 173-194.

12. Rigel DS, Wheeland RG. Deaths related to liposuction (letter; comment). *N Engl J Med* 1999;341:1001-1002; discussion:1002-1003.

13. Shiffman M. Medications potentially causing lidocaine toxicity. *American J Cosmetic Surg* 1998;15:227-228.

14. Klein JA. Superwet liposuction and pulmonary edema. In: Tumescent technique tumescent anesthesia and microcannular liposuction. St. Louis: Mosby; 2000. p. 61-66.

15. Lillis PJ. Liposuction surgery under local anesthesia: limited blood loss and minimal lidocaine absorption. *J Dermatolog Surg Oncology* 1988;114:1145-1148.

16. Rohrich RJ, Beran SJ. Is liposuction safe? Plast Reconstr Surg 1999;104;3:819-822.

17. Hanke CW, Bernstein G, Bullock BS. Safety of tumescent liposuction in 15,336 patients „ national survey results. *Dermatolog Surg* 196;22:459-462.

Bibliography

American Society for Dermatological Surgery: Liposuction Guidelines 200;26:265-269.

Brain SD, Williams TJ. Inflammatory oedema induced by synergism between calcitonin gene-related peptide (CGRP) and mediators of increased vascular permeability. *Br J Anesth* 1985;86:855-860.

Breuninger H, Hobbach P, Schimek F. Ropivacaine: an important anesthetic agent for slow infusion and other forms of tumescent anesthesia. *Dermatologic Surgery* 1999;25:799-802.

Coleman WP III, Katz B, Bruck M, Narins R, Lawrence N, Flynn TC, et al. The ef"cacy of powered liposuction. *Dermatolog Surg* 2001;27:735-738.

Deeb M, Eed A. Megaliposuction: analysis of 1520 patients. *Aesthetic Plast Surg* 1999;123:16-22.

Di Giuseppe A. Ultrasonically assisted liposculpturing: physical and technical principles and clinical applications. *Am J Cosmetic Surg* 1997;14:317-327.

Di Giuseppe A. Ultrasound-assisted liposuction: physical and technical principles. In: Shiffman, MA, Di Giuseppe A, editors. Liposuction, principles and practice. New York: Springer-Verlag; 2006. p. 229-238.

Esclamado RM, Damiano GA, Cummings CW. Effect of local hypothermia on early wound repair. *Arch Otolaryngol Head Neck Surg* 1990;116:803-808.

Fischer G. First surgical treatment for modeling body•s cellulite with three 5 mm incisions. *Bull Int Acad Cosm Surg* 1976;2:35-37.

Illouz YG. Body contouring by lipolysis: a 5-year experience with over 3000 cases. *Plast Reconstr Surg* 1983;72:591-597.

Kaminski MV, Lopez de Vaughan RM. The anatomy and physiology metabolism / nutrition of subcutaneous fat. In: Shiffman MA, Di Giuseppe A, editors. Liposuction, principles and practice. New York: Springer-Verlag; 2000. p. 17-24.

Klein JA. Tumescent liposuction with local anesthesia. In: Lask GP, Moy RL, editors. Principles and techniques of cutaneous surgery. New York: McGraw Hill; 1996. p. 529-542.

Parish TD. Pharmacokinetics of tumescent anesthesia. In: Shiffman, M.A., Di Giusseppe A, editors. Liposuction, Principles and Practice. New York: Springer-Verlag; 2006. p. 17-24.

Pitman GH, Teimouran B. Suction lipectomy: complications and results by survey. *Plast Reconstr Surg* 1985;76:65-72.

Rohrich RJ, Beran SJ, Fodor FB. The role of subcutaneous in"ltration in suction-assisted lipoplasty: a review. *Plast Reconstr Surg* 1997;99:514-519.

Salans LB, Cushman SW, Weismann RE. Studies of human adipose tissue, adipose cell size and number in nonobese and obese patients. *J Clin Invest* 1973;52:929-941.

Scott DB. Evaluation of clinical tolerance of local anesthetic agents. *Br J Anesth* 1975;47:328-333.

Shiffman M.A. Principles of liposuction. In: Shiffman MA, Di Giuseppe A, editors. Liposuction, principles and practice. New York: Springer-Verlag; 2006. p. 379-380.

Sommer B. Vorteile und Nachteile der Tumeszenz-Lokalanastehesie. In: Sommer B, Sattier G, Hanke CW, editors. Tumeszenz-Lokalanastehesie. Berlin: Springer-Verlag; 1998. p. 40-44.

Reprint request: Daniel Man, MD, 851 Meadows Road, Suite 222, Boca Raton, FL 33486.

Copyright © 2007 by The American Society for Aesthetic Plastic Surgery, Inc.

1090-820X / $32.00

doi:10.1016.j.asj.2007.04.008